

Table Nitidulid pests of stored products introduced in Europe and Mediterranean areas (After Jelinek et al. 2016)

Species	Origin	Location and Year Introduced
<i>Carpophilus binestratus</i>	Palaeotropical	Mediterranean areas; probably 1700 (may be within original range)
<i>Carpophilus dimidiatus</i>	Palaeotropical	Mediterranean areas; probably 1700 (mostly dried fruit)
<i>Carpophilus fumatus</i>	Afrotropical	Portugal (Azores, 1985; Madeira, 1996); Cape Verde Islands, 1988; Italy, 1990; Madeira 1996; Near East, Morocco 2000
<i>Carpophilus hemipterus</i>	Paleotropical	Mediterranean areas; probably Roman Age
<i>Carpophilus ligneus</i>	Central America	England 1940; Greece; Northern Africa; Canary Islands; Germany
<i>Carpophilus lugubris</i>	Southern USA	NE Italy (cornfields, mature strawberries)
<i>Carpophilus marginellus</i>	Paleotropical	Mediterranean areas; Europe, including Fennoscandia and Ireland; probably 1800/1900
<i>Carpophilus mutilatus</i>	Paleotropical	Mediterranean areas; probably 1800/1900
<i>Carpophilus nepos</i> , syn. <i>C. freemani</i>	Paleotropical	Mediterranean areas; probably 1800/1900
<i>Carpophilus obsoletus</i>	Paleotropical	Mediterranean areas and Belgium, probably 1700
<i>Carpophilus succisus</i>	Neotropical	Portugal (Azores, 2000)
<i>Carpophilus truncatus</i>	Paleotropical	Mediterranean areas; probably 1800 (mostly dried fruit)
<i>Carpophilus zeaphilus</i>	East Africa	Portugal, 1985; Spain; Monaco; Italy (Sicily, peninsular Italy); Albania; Near East, 1991/2015
<i>Epuraea luteolus</i>	Oriental	Israel and Near East, 1970; Mediterranean and southern European areas, 1980
<i>Epuraea ocularis</i>	Oriental	Spain (Canary Islands, 1992); Portugal (Madeira, 1996); south and central Europe; Moldova; Turkey; Caucasus
<i>Glischrochilus fasciatus</i>	USA	Switzerland, 1992; Germany; Austria; Slovenia; Finland
<i>G. quadrisignatus</i>	USA	Germany, 1945; Italy, 1985, central and SE Europe

<i>Lobiopa insularis</i>	Tropical South and Central America	Spain (Canary Islands; Tenerife, 2008) (dates)
<i>Omosita funesta</i>	Mexico	Spain (Teruel, 1931) (probably imported sausage)
<i>Phenolia tibialis</i>	west tropical Africa	Azores and Madeira, 1985; continental Portugal, 2008; Spain
<i>Stelidota geminata</i>	Southern USA	Portugal (Azores Islands, Flores, 1984); northern Italy (Lombardy, Sicily, 1996/1998); Belgium; Spain (including Canary Islands); France; Switzerland; Slovenia; Austria; Germany; Hungary; Turkey; Russian Caucasus, 1993/2015
<i>Urophorus humeralis</i> syn. <i>Carpophilus humeralis</i>	Paleotropical	Europe, probably 1900
